

One pot MCR for the 9-phenyl-1,2,3,4,5,6,7,8,9,10-decahydroacridine derivatives for good yield via Silica supported green catalyst and solvent free condition

JAYAVEERSINH MAHIDA, RAVI B. PATEL

Shree P. M. Patel Institute of P.G. Studies & Research in Science, Anand, Gujarat, India

Abstract: Organic synthesis is the very useful for the modification of the drug discovery and its applicable for the modify for the change the route of drug discovery as well as pharmacology study, the current situation of the environment is very harsh and environmentally friendly sources are costly then the tradition method of synthesis, so our worked is focus on designing the environment as well as economically viable route of synthesis with the help of some novel silica supported metal catalyst. in this type catalyst design from collected solid harmful environment metal waste of aluminum foil. our team is collected aluminum waste from food sellers and then wash with water aluminum converted to aluminum hydroxide. they use for the formulation of gel formation of medicine. and its not harmful like aluminum metal, as per survey aluminum is the second highest metal waste on the earth.

Keywords: Dimidone, Solvent free, green catalyst. high reusability.



1. INTRODUCTION

This catalyst are design for the homogeneous catalyst for the differ path for the new heterogeneous catalyst. Aluminum hydroxide is the base its docking on the porous silica. for the increase the activity of the catalyst with lowest amount of the catalyst. the nano particle, Green catalyst and increase the reusability or activity of catalyst is the immerging field for the new catalyst design and change the reaction route for the deployment in the synthesis field, so our team is focused on this thing, our goal is the docking molecule on silica, green catalyst and its reusability is almost double then other nano particle or docking material they used as catalyst in previous work. design the catalyst motivated from Nasser H et al, they sodium aluminum silicates for the abrogation of dyes from environment, and Ahemad nakhaei et al

they are introduce the heterogeneous catalyst for the 1,4 dihydropyridine Hantzsch reaction for modification study, Aluminum, ferrous, Titanium like metals nano particle or its compositions are used as a catalyst in reactions, they required 30 to 460 min for the ending of reaction. and they reusability is approximately 5 cycles they are gave good yield after the 5 cycles they continue reduced its yield. they are gave not effective yield in case silica supported results but our team are introduce successful silica supported aluminum hydroxide have good activity and reusability for this type reactions.

Material and method:

Silica supported $\text{Al}(\text{OH})_3$: First collect the waste aluminum foil from food court then wash with water and remove all stacked material of food, then cuts in small pieces after cutting the dissolve in waste HCl they HCl is by product of pharmaceutical company, the addition of aluminum pieces is slowly in HCl at the time of the addition the evaluation of hydrogen gas and rise the temperature and then the cool it, after the cool the addition of Sodium hydroxide pallet and set $\text{pH}=2.0$ to 7.0 and above and white precipitation is formed, its filter by vacuum filter and dry in hot air oven for 90°C for 2 hour after the drying the make the slurry of Silica and the addition of aluminum hydroxide in slurry and starring for 2 hours after the remove the solvent from slurry the dry and kept in muffle furnace for 1 hour 300°C after this the fine powder is ready they used as a green catalyst, it's conform by SEM, XRD, BET analysis.



Experimental section: 2.5 mmole aldehyde, 5 mmole cyclohexane-1,3-dione with 1.5 mmole Ammonium acetate mix well then addition of catalyst in variety of amount with different temperature and check the efficiency of catalyst. monitor the reaction on TLC plate after the completion of reaction the all reaction mass dissolve in suitable solvent like MeOH, EtOH at the time of purification process all product are dissolve in solvent then filter it and collect the catalyst and wash with EtOH and use for the next cycle.

Result and discussion: this synthesis is focused on the green as well as novel silica supported catalyst for the economically viable synthesis route for the deployment of new thing in catalysis. the good yield, high reusability, so, our team are waste aluminum foil converted to efficient and capable green catalyst for the env-eco catalyst. they are gave the same efficiency other catalyst like SbCl_3 , HPA1, HPA2, LGPCO, the check appropriate time, temperature and amount of the catalyst for this reaction, we conclude that the solvent free condition and the various solvent are use for the reaction at variety of temperature for the optimization of reactions condition for the good yield, and after the all solvent.

Table:01 comparative study with previously reported catalyst

Sr	Amount of catalyst	Solvent	Temp ($^{\circ}\text{C}$)	Time	Yield(%)	Yield (Rp)
1	$\text{Al}(\text{OH})_3(0.1)$	Solvent free	110°C	15-30	64-75 ^(TW)	90 ⁽²⁷⁾
2	$\text{Al}(\text{OH})_3(0.2)$	Solvent free	110°C	15-30	67-77 ^(TW)	NR
3	$\text{Al}(\text{OH})_3(0.3)$	Solvent free	110°C	15-30	71-79 ^(TW)	94 ⁽³¹⁾
4	$\text{Al}(\text{OH})_3(0.4)$	Solvent free	110°C	15-30	73-82 ^(TW)	96 ⁽³¹⁾
5	$\text{Al}(\text{OH})_3(0.5)$	Solvent free	110°C	15-30	73-84 ^(TW)	93 ⁽³¹⁾
6	Sio- $\text{Al}(\text{OH})_3(0.1)$	Solvent free	110°C	15-30	71-74 ^(TW)	94 ⁽³¹⁾
7	Sio- $\text{Al}(\text{OH})_3(0.2)$	Solvent free	110°C	15-30	72-81 ^(TW)	89 ⁽³²⁾
8	Sio- $\text{Al}(\text{OH})_3(0.3)$	Solvent free	110°C	15-30	75-82 ^(TW)	NR
9	Sio- $\text{Al}(\text{OH})_3(0.4)$	Solvent free	110°C	15-30	79-89 ^(TW)	95 ⁽³¹⁾
10	Sio- $\text{Al}(\text{OH})_3(0.5)$	Solvent free	110°C	15-30	91-96 ^(TW)	
11	$\text{Y}(\text{otf})_3(5\%)$	H_2O	100°C	360-420	360-420	87 ⁽³⁾
12	HCl	H_2O	100°C	360-420	360-420	49 ⁽⁴⁾
13	TFA	H_2O	100°C	360-420	360-420	55 ⁽⁵⁾
14	HoAC	H_2O	100°C	360-420	360-420	24,25,29,32,31 ⁽⁴⁾
15	$\text{SbCl}_3\text{-SiO}_2$	Solvent free	120°C	50	91 ⁽⁶⁾	

16	$\text{Zn}(\text{OAc})_2$	H_2O	120
17	Any Organic acid	H_2O	120
18	HPA1, HPA2	Solvent free	110
19	$\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$	IL	110
20	TBA	Solvent free	100
21	CBSA	Solvent free	100
22	$\text{B}(\text{C}_6\text{F}_5)_3$	Solvent free	R
23	SBSSA	EtOH	65
24	Amberlyst-15	CH_3CN	Ref
25	LGPCO	Solvent free	Ref
26	GSMO	Solvent free	Ref
27	$\text{ZnCl}_2\text{-SiO}_2$	Solvent free	Ref
28	$\text{FeCl}_3\text{-SiO}_2$	Solvent free	Ref

Table:02

time:15-30 min, Amount of catalyst:0.5 mmole, temp:110 $^{\circ}\text{C}$, NF-Not formed, NR-Not reported

*PC	Aldehyde	Yield (TW)	Yield (Rp)
JBR#01	C_6H_5	96	90 ⁽²⁷⁾
JBR#02	2- NO_2 C_6H_5	NF	NR
JBR#03	3- NO_2 C_6H_5	96	94 ⁽³¹⁾
JBR#04	4- NO_2 C_6H_5	99	96 ⁽³¹⁾
JBR#05	2- Cl C_6H_5	96	74 ⁽³²⁾
JBR#06	4- Cl C_6H_5	92	88 ⁽³¹⁾
JBR#07	3- Cl C_6H_5	94	93 ⁽³¹⁾
JBR#08	4- Cl C_6H_5	97	94 ⁽³¹⁾
JBR#09	3,4- Br - MeOC_6H_3	93	89 ⁽³²⁾
JBR#10	3- Br C_6H_5	NF	NR
JBR#11	4- Br C_6H_5	97	95 ⁽³¹⁾

Conclusion: our team are selected the good yield contained derivatives for the study of catalysis potential for the reaction via green way, we got the highest amount of yield at the 0.5% at the 110°C and got the 91-96% of yield the past researchers formed 90-96% (3,4,5,6,360,420,24,25,29,32,31), but the High time 50 the completion of the reactions. so, we are tray to optimization of the waste

after the reaction for the and we go the highest amount of yield in very short time and the more productive thing of this catalyst the recrystallization does not required they are the complete conversion of the reaction but the spectroscopic conformation we are the performed the recrystallization on via hot Analytical data:

JBR#01

9-phenyl-3,4,6,7,9,10-hexahydroacridine-1,8(2H,5H)-dione:

IR:1226(-C-Nstr),1487,1529(CH₂-CH₂),1635(-C=O,Carbonylgroup),2960(CHstr),3054(CH Str),3755(NH str)

NMR:69.02(1to5ppm,12.43=12H,allnon aromatic ring hydrogen),28.02(6to 8,5.088=5H,aromatic ring hydrogen),3.57(0.64=1H,nitrogen contenting ring hydrogen)

¹³CNMR:28.77,30.66,44.43,52.82,119.35,119.55,119.99,123.62,124.34,129.02,129.10,,137.46,138.45,138.56,147.24,149.26,204.78.

Mass:293.98

JBR#04

9-(4-nitrophenyl)-3,4,6,7,9,10-hexahydroacridine-1,8(2H,5H)-dione:

IR: 1231,1342,1490,1608,1659,2958,3060,3172

¹HNMR:71.85(1to5ppm,13.65=13H,all non aromatic ring hydrogen),28.02(6to 8,5.088=5H,aromatic ring hydrogen),3.57(0.64=1H,nitrogen contenting ring hydrogen)

¹³CNMR:23.20,31.50,33.42,37.90,114.31,124.78,127.25,128.44,131.52,148.78,153.20

MASS:339.25

JBR#08

9-(4-chlorophenyl)-3,4,6,7,9,10-hexahydroacridine-1,8(2H,5H)-dione

¹HNMR:74.70(1to5ppm,13.45=13H,all non aromatic ring hydrogen),20.42(6to 8,3

.62 =4H,aromatic ring hydrogen),6.21(1.11=1H,nitrogen contenting ring hydrogen)

¹³CNMR:23.32,30.44,33.78,36.39,42.70,111.18,115.55,111.78,1129.44,130.15,131.78,142.58,155.47,162.21,200.45

Mass:327.10

ethanol. the variety of derivatives are the formed by the change the substitution of aromatic aldehyde and we got the different derivatives of hexahydroacridine. the conformation of the structure with help of IR by FT-IR Proton NMR, Carbon NMR and Mass spectra.

References:

- [1]Sun,T.,Li,H.,Wu Z.,Li C.,Ren, X.,and Zhang, F.(2018) Three-component Strategy for Synthesis of Dihydropyrido2,3-d.pyrimidine and Spirooxindole Derivatives. *J. Heterocyclic Chem.*, 55: 2270–2276.
- [2] Mabbett, A. N., Sanyahumbi, D., Yong, P., & Macaskie, L. E. (2006). Biorecovered precious metals from industrial wastes: single-step conversion of a mixed metal liquid waste to a bioinorganic catalyst with environmental application. *Environmental science & technology*, 40(3), 1015-1021.
- [3]Deplanche, K., Merroun, M. L., Casadesus, M., Tran, D. T., Mikheenko, I. P., Bennett, J. A., ... & Selenska-Pobell, S. (2012). Microbial synthesis of core/shell gold/palladium nanoparticles for applications in green chemistry. *Journal of the Royal Society Interface*, 9(72), 1705-1712.
- [4]Dang, G. H., Vu, Y. T., Dong, Q. A., Le, D. T., Truong, T., & Phan, N. T. (2015). Quinoxaline synthesis via oxidative cyclization reaction using metal–organic framework Cu (BDC) as an efficient heterogeneous catalyst. *Applied Catalysis A: General*, 491, 189-19.
- [5] Darabi, H. R., Aghapoor, K., Mohsenzadeh, F., Taala, F., Asadollahnejad, N., & Badiei, A. (2009). Silica-supported antimony (III) chloride as highly effective and reusable heterogeneous catalyst for the synthesis of quinoxalines. *Catalysis letters*, 133(1-2), 84.
- [6] Bagley, M. C., Dale, J. W., & Bower, J. (2002). A new one-pot three-component condensation reaction for the synthesis of 2, 3, 4, 6-tetrasubstituted pyridines. *Chemical Communications*, (16), 1682-1683.
- [7] Aghapoor, K., Ebadi-Nia, L., Mohsenzadeh, F., Morad, M. M., Balavar, Y., & Darabi, H. R. (2012). Silica-supported bismuth (III) chloride as a new recyclable heterogeneous catalyst for the Paal–Knorr pyrrole synthesis. *Journal of Organometallic Chemistry*, 708, 25-30.
- [8] Kadam, H. K., Khan, S., Kunkalkar, R. A., & Tilve, S. G. (2013). Graphite catalyzed green synthesis of quinoxalines. *Tetrahedron Letters*, 54(8), 1003-1007.
- [9] Nageswar, Y. V. D., Reddy, K. H. V., Ramesh, K., & Murthy, S. N. (2013). Recent developments in the synthesis of quinoxaline derivatives by green synthetic approaches. *Organic Preparations and Procedures International*, 45(1), 1-27.
- [10]Sabitha,G.;Reddy,G.K.K.;Reddy,C.S.;Yadav,J. S.*Tetrahedron Lett.* **2003**, 44, 4129.

- [11] Lei, M.; Ma, L.; Hu, L. *Synth. Commun.* **2011**, *41*, 1969.
- [12.] Mousavi, M.R.; Maghsoodlou, M.T.; Hazeri, N.; Aboonajmi, J.; Habibi-Khorassani, S.M. *Iran. J. Org. Chem.* **2016**, *6*, 1323
- [13] Wang, X.; Gong, H.; Quan, Z.; Li, L.; Ye, H. *Synth. Commun.* **2011**, *41*, 3251.
- [14] Datta, B.; Pasha, M.A. *Chin. J. Catal.* **2011**, *32*, 1180.
- [15] Sabitha, G.; Arundhati, K.; Sudhakar, K.; Sastry, B.S.; Yadav, J.S. *Synth. Commun.* **2009**, *39*, 2843.
- [16] Khodja, I.A.; Ghalem, W.; Dehimat, Z.I.; Boulcina, R.; Carboni, B.; Debache, A. *Synth. Commun.* **2014**, *44*, 959.
- [17]. Zolfigol, M.A.; Salehi, P.; Khorramabadi-Zad, A.; Shayegh, M. *J. Mol. Catal. A: Chem.* **2007**, *261*, 88.
- [18]. Debache, A.; Ghalem, W.; Boulcina, R.; Belfaitah, A.; Rhouati, S.; Carboni, B. *Tetrahedron Lett.* **2009**, *50*, 5248.
- [19] Debache, A.; Chouguiat, L.; Boulcina, R.; Carboni, B. *Open Org. Chem. J.* **2012**, *6*, 12.
- [20] Murthy, Y.L.N.; Rajack, A.; Ramji, M.T.; Praveen, C.; Lakshmi, K.A. *Med. Chem. Lett.* **2012**, *22*, 6016.
- [21] Shockravi, A.; Kamali, M.; Sharifi, N.; Nategholeslam, M.; Moghanlo, S.P. *Synth. Commun.* **2013**, *43*, 1477.
- [22] Tewari, N.; Dwivedi, N.; Tripathi, R.P. *Tetrahedron Lett.* **2004**, *45*, 9011.
- [23] Jahanbin, B.; Davoodnia, A.; Behmadi, H.; Tavakoli-Hoseini, N. *Bull. Korean Chem. Soc.* **2012**, *33*, 2140.
- [24] AhmadNakhaei*, AbolghasemDavoodnia, SepidehYadegarian, NiloofarTavakoli-Hoseini. *Iranian Journal of Organic Chemistry* Vol. 8, No. 4 (2016) 1919-1927.
- [25] Smita T. Morbale, Sachin K. Shinde, Shashikant A. Damate, Madhukar B. Deshmukhand Suresh S. Patil. *Letters in Organic Chemistry*, 2018, 15, 57-63.
- [26] Ahooie, T.S., Azizi, N., Yavari, I. et al. *J IRAN CHEM SOC* (2018) 1 5: 855.
- [27] B.27, H. Eshghi, A. Khojastehnezhad, R. Tayebee S. Sedigh Ashraf, G. Esmailian Kahoo, F. Moeinpour, *RSC Adv.* **5**, 64850 (2015).
- [28] N Azizi, N. Dado, A. K Amiri, *Can. J. Chem.* **90**, 195 (2012).
- [29] Najmaddin Azizi, Nairreh Dado, and Alireza Khajeh Amiri, *Can. J. Chem.* **90**, 2, 167, 2012.
- [30] M. Esmailpour, J. Javidi, F. Dehghani, S. Zahmatkesh, *Res. Chem. Intermed.* **43**, 163 (2017)
- [31] Tavakkoli, H., Sanaeishoar, H., Mohave, F. et al. *Reac Kinet Mech Cat* (2016) 119: 259.
- [32] Ramesh, K. B., & Pasha, M. A. (2014). Study on one-pot four-component synthesis of 9-aryl-hexahydro-acridine-1, 8-diones using SiO₂-I as a new heterogeneous catalyst and their anticancer activity. *Bioorganic & medicinal chemistry letters*, *24*(16), 3907-3913.

3. MAIN TITLE

The main title (on the first page) should begin 1 3/16 inches (7 picas) from the top edge of the page, centered, and in Times New Roman 14-point, boldface type. Capitalize the first letter of nouns, pronouns, verbs, adjectives, and adverbs; do not capitalize articles, coordinate conjunctions, or prepositions (unless the title begins with such a word) (*e.g.* **A Comparative Study of Privacy Protection Methods for Smart Home Environments**). Please initially capitalize only the first word in other titles, including section titles and first, second, and third-order headings (for example, “Titles and headings” — as in these guidelines). Leave two blank lines after the title.

4. Author Name(s) and Affiliation(s)

Author names and affiliations are to be centered beneath the title and printed in Times New Roman 12-point, non-boldface type. (See example below)

4.1. Affiliations

Affiliations are centered, italicized, not bold. Include e-mail addresses if possible.

For example:

Author¹, Author² and Author³

¹*Affiliation*

²*Affiliation*

³*Affiliation*

¹*Email*, ²*Email*, ³*Email*

4.2. Corresponding Author

Corresponding author should have an asterisk sign (*) if possible, after the corresponding author's name. The Corresponding author (e.g., *Corresponding Author) label should be appeared at the footnote section of the first page of the paper, Times New Roman in style and 10 in font size.

5. Second and Following Pages

The second and following pages should begin 1.0 inch (2.54 cm) from the top edge. On all pages, the bottom margin should be 1-3/16 inches (2.86 cm) from the bottom edge of the page for 8.5 x 11-inch paper; for A4 paper, approximately 1-5/8 inches (4.13 cm) from the bottom edge of the page.

6. Type-style and Fonts

Wherever Times New Roman is specified, Times New Roman may be used. If not available in your word processor, please use a font closest to Times New Roman that you have access to. Please avoid using bit-mapped fonts if possible. True-Type 1 fonts are preferred.

7. Main Text

Type your main text in 11-point Times New Roman, single-spaced. Do not use double-spacing. All paragraphs should be indented 1 pica (approximately 1/6- or 0.17-inch or 0.43 cm). Be sure your text is fully justified, flush left and flush right. Please do not place any additional blank lines between paragraphs.

7.1. Tables

Place tables as close as possible to the text they refer to and aligned center. A table is labeled *Table* and given a number (e.g., **Table 1. Sample Datasheet with Attributes in Linguistic Term**) it should be numbered consecutively. The table label and caption or title appears 9pt space above the table, 6pt space after the text or paragraph if any; it should be uniform fonts and font size, and use 11pt font size and Helvetica style, capitalized similar to paper title, aligned center and bold face. Sources and notes appear below the table, aligned left. All tables must be in portrait orientation.


For Example:

Table 1. Table Label

7.2. Figures

Place figures as close as possible to the text they refer to and aligned center. Photos, graphs, charts or diagram should be labeled *Figure* (do not abbreviate) and appear 6pt space below the figure, 12pt space before the next text or paragraph, and assigned a number consecutively. The label and title should be in line with the figure number (*e.g.*, **Figure 1. Location Error Rate of Three Schemes**), it should be uniform fonts and font size; use 11pt font size and Helvetica style, capitalized similar to paper title, aligned center and bold face. Source (if any) appear underneath, flush left. Figures should be at good enough quality. Minimum image dimensions are 6 cm (2.3622 in) wide by 6 cm (2.3622 in) high.

For Example:

**Figure 1. Figure Label**

7.3. Equations

Including symbols and equations in the text, the variable name and style must be consistent with those in the equations. Equations should be indented at the left margin and numbered at the right margin, equation number is enclosed with open and close parenthesis () Time New Roman in style and 11pt font size. Define all symbols the first time they are used. All equation symbols must be defined in a clear and understandable way.

For Example:

$$\varphi_{\mu\nu}(z) = \frac{\|k_{\mu\nu}\|^2}{\sigma^2} e^{-\frac{\|k_{\mu\nu}\|^2 \|z\|^2}{\sigma^2}} [e^{ik_{\mu\nu}z} - e^{-\frac{\sigma^2}{2}}] \quad (1)$$

8. First-order Headings

For example, “**1. Introduction**”, should be Times New Roman 13-point boldface, initially capitalized, flush left, with one blank line before, and one blank line after.

8.1. Second-order Headings (Sub-heading)

As in this heading, they should be Times New Roman 11-point boldface, initially capitalized, flush left, with one blank line before, and one after.

8.1.1. Third-order Headings: Third-order headings, as in this paragraph, are discouraged. However, if you must use them, use 11-point Times New Roman, boldface, initially capitalized, flush left, and preceded by one blank line, followed by a colon and your text on the same line.

9. Footnotes

Use footnotes sparingly (or not at all) and place them at the bottom of the column of the page on which they are referenced to. Use Times New Roman 9-point type, single-spaced. To help your readers, avoid using footnotes altogether and include necessary peripheral observations in the text (within parentheses, if you prefer, as in this sentence).

Appendix

An appendix, if needed, should appear before the acknowledgments.

Acknowledgments

These should be brief and placed at the end of the text before the references.

REFERENCES

List and number all bibliographical references that has important contribution on the paper, (if possible, limit to 30, which only are necessary citations are recommended). 9-point Times New Roman, fully justified, single-spaced, at the end of your paper. When referenced in the text, enclose the citation number in square brackets, for example [1]. Do not abbreviate the months. Don't forget to put period (.) at the end of each reference. (See examples below)

11.1. Journal Article

- [1] C. D. Scott and R. E. Smalley, “Diagnostic Ultrasound: Principles and Instruments”, *Journal of Nanosci. Nanotechnology.*, vol. 3, no. 2, (2003), pp. 75-80.

11.2. Book

- [2] H. S. Nalwa, Editor, “Magnetic Nanostructures”, American Scientific Publishers, Los Angeles, (2003).

11.3. Chapter in a Book

- [3] H. V. Jansen, N. R. Tas and J. W. Berenschot, “Encyclopedia of Nanoscience and Nanotechnology”, Edited H. S. Nalwa, American Scientific Publishers, Los Angeles, vol. 5, (2004), pp. 163-275.

11.4. Conference Proceedings

- [4] *J. Kimura and H. Shibasaki, "Recent Advances in Clinical Neurophysiology", Proceedings of the 10th International Congress of EMG and Clinical Neurophysiology, Kyoto, Japan, (1995) October 15-19.*

11.5. Patent

- [5] *C. E. Larsen, R. Trip and C. R. Johnson, "Methods for procedures related to the electrophysiology of the heart", U.S. Patent 5,529,067, (1995) June 25.*